

ON THE REPRESENTATION DIMENSION OF RANK 2 GROUP ALGEBRAS AND RELATED ALGEBRAS

THORSTEN HOLM
University of Leeds

WEI HU
University of Leeds/
Beijing Normal University

Abstract

The representation dimension was defined by M. Auslander in 1970 and is, due to spectacular recent progress, one of the most interesting homological invariants in representation theory. The precise value is not known in general, and is very hard to compute even for small examples. For group algebras, it is known in the case of cyclic Sylow subgroups, due to Auslander's fundamental work. For some group algebras (in characteristic 2) of rank at least 3 the precise value of the representation dimension follows from recent work of R. Rouquier. There is a gap for group algebras of rank 2; here the deep geometric methods do not work. In this paper we show that for all $n \geq 0$ and any field k the commutative algebras $k[x, y]/(x^2, y^{2+n})$ have representation dimension 3. For the proof, we give an explicit inductive construction of a suitable generator-cogenerator. As a consequence, we obtain that the group algebras in characteristic 2 of the groups $C_2 \times C_{2^m}$ have representation dimension 3. Note that for $m \geq 3$ these group algebras have wild representation type.

MSC Classification: 16G10 (primary); 13D05, 16S50, 20C05 (secondary).

1 Introduction

The representation dimension of an Artin algebra was defined by M. Auslander [1] as a way of measuring homologically how far an algebra is from being of finite representation type. (For the precise definition see Section 2 below.) In fact, Auslander showed that the representation dimension of an algebra is at most 2 if and only if the algebra has finite representation type, i.e. has only finitely many indecomposable modules (up to isomorphism).

For more than two decades it remained unclear whether Auslander's philosophy and hopes were justified, one of the main problems being that the precise value of the representation dimension of an algebra is very hard to determine. For instance, until the late 1990's there were still only very few examples of

algebras known with representation dimension 3, and not a single algebra with representation dimension > 3 .

This situation changed dramatically over the last few years, with spectacular recent progress which we briefly mention here. O. Iyama proved that the representation dimension is always finite [6]. This integer attached to any algebra is actually an invariant under various notions and levels of equivalences of algebras. As it is defined, the representation dimension is invariant under Morita equivalences. In modern representation theory the notions of stable equivalences and especially derived equivalences became increasingly important. It was recently shown by X. Guo [4] that the representation dimension is invariant under stable equivalences. (For stable equivalences of Morita type this was already shown by C. Xi [10].) In particular, for selfinjective algebras the representation dimension is invariant under derived equivalences, thus providing one of the few explicit invariants of derived module categories known so far.

A breakthrough was obtained by R. Rouquier [8] (but see also the new revised version [9]). As a consequence of a deep general theory of dimensions of triangulated categories he obtained the first examples of algebras with representation dimension > 3 , solving one of the most fundamental open questions about this invariant. Much more than that, Rouquier obtained that the representation dimension is unbounded, by showing that for any $n \geq 3$ the exterior algebra of an n -dimensional vector space has representation dimension $n + 1$. Actually, the methods used there indicate a very geometric nature of the representation dimension in these cases, and from this approach one can expect many more explicit examples in the near future.

These recent results finally make it clear that Auslander's philosophy of measuring how far an algebra is from being of finite representation type is working; we get a new division of algebras according to the size of their representation dimension. Note that this new division certainly does not run along the lines of the classical division into finite, tame and wild representation type. For instance, already Auslander showed that all finite-dimensional hereditary algebras have representation dimension at most 3, not depending on their representation type. There is something completely new to be discovered! So far, it remains mysterious in general what structural properties of an algebra actually determine the representation dimension.

In this paper, we will provide more examples of algebras of wild representation type having representation dimension 3. Our main motivation comes from group algebras. It is well known that the representation type of a modular group algebra kG of a finite group over a field k of characteristic p is determined by a Sylow p -subgroup P of G [2]: kG has finite representation type if and only if P is cyclic; kG has tame representation type precisely when $p = 2$ and P is dihedral, semidihedral or quaternion. In all other cases kG has wild representation type. Hence, in the cyclic case, the representation dimension of kG is at most 2. It is also known that all group algebras of tame representation type (and related algebras) have representation dimension 3. (Unfortunately, there is no complete proof in the literature, but see [5], and [7], 4.4.2.)

From Rouquier's work one gets that for any $n \geq 3$ the mod 2 group algebra

of an elementary abelian group $C_2 \times \dots \times C_2$ of order 2^n has representation dimension $n + 1$. One could hope that the geometric methods used could be adapted and generalized to deal with elementary abelian p -groups in odd characteristic as well. However, these geometric methods do not work for elementary abelian groups of rank 2.

The main aim of this paper is to give some partial results on the rank 2 case, thus providing the first examples to fill this 'rank-2-gap'. Note that the mod p group algebra of $C_{p^m} \times C_{p^n}$ is isomorphic to the commutative ring $k[x, y]/(x^{p^m}, y^{p^n})$. So the natural class of algebras to study are the commutative rings $A_{n,m} := k[x, y]/(x^m, y^n)$, for natural numbers $n, m \geq 2$.

Our main result in this paper gives the precise value of the representation dimension for the algebras $A_{2,m}$. Note that this result is independent of the ground field.

Theorem 1.1 *Let k be a field and let $\Lambda_n = k[x, y]/(x^2, y^{2+n})$ for every non-negative integer n . Then the representation dimension of Λ_n is 3.*

As a consequence we get the following partial answer on the 'rank-2-gap' for group algebras.

Corollary 1.2 *Let k be a field of characteristic 2. Then for any $n \geq 1$ the group algebra $k(C_2 \times C_{2^n})$ has representation dimension 3.*

The paper is organized as follows. In Section 2 we recall the necessary definitions and background on representation dimensions. In particular, we outline the method for determining the precise value of the representation dimension. The final Section 3 then contains the proof of the main result.

2 Computing representation dimensions

The representation dimension has been defined by M. Auslander [1] for arbitrary Artin algebras. However, for the purpose of this paper, by an algebra we mean a finite-dimensional algebra over a fixed field k . (It will turn out that our results will not depend on the ground field.)

A finitely generated A -module M is called a *generator-cogenerator* for A if all projective indecomposable modules and all injective indecomposable A -modules occur as direct summands of M .

Auslander's basic idea was to study homological properties of endomorphism rings of modules (instead of studying modules directly). For any algebra A the *representation dimension* is defined to be

$$\text{repdim}(A) := \inf\{\text{gl.dim}(\text{End}_A(M)) \mid M \text{ generator-cogenerator}\}.$$

Note that this invariant a priori has values in $\mathbb{N}_0 \cup \{\infty\}$. However, it was shown by O. Iyama [6] that $\text{repdim}(A)$ is always finite.

Auslander's fundamental result states that $\text{repdim}(A) \leq 2$ if and only if A has finite representation type. More precisely, $\text{repdim}(A) = 0$ if and only if A is semisimple, and $\text{repdim}(A) \neq 1$ for all algebras A .

The definition indicates how to determine upper bounds for $\text{repdim}(A)$; in fact, if one can show for some generator-cogenerator M that $\text{gl.dim}(\text{End}_A(M)) \leq m$ then $\text{repdim}(A) \leq m$. In particular, if one can find a generator-cogenerator M with $\text{gl.dim}(\text{End}_A(M)) \leq 3$, and if A is not of finite representation type, then one has shown that $\text{repdim}(A) = 3$. Of course, the notoriously difficult problem is to find a suitable module M .

However, this is exactly the strategy we will pursue successfully in the proof of our main theorem. We will explain in Section 3 below how to construct inductively generator-cogenerators for the algebras $k[x, y]/(x^2, y^{2+n})$ with endomorphism rings of global dimension 3.

We now describe the general method used later for how to determine the global dimension of the endomorphism ring of a generator-cogenerator M .

Let M be a finitely generated module over a finite-dimensional algebra A . The identity of $\text{End}_A(M)$ is the sum of the "identity maps" on the indecomposable direct summands of M . Hence we have primitive idempotents of $\text{End}_A(M)$ corresponding to the summands of M . For any indecomposable summand T of M we denote the corresponding simple $\text{End}_A(M)$ -module by E_T . The corresponding indecomposable projective $\text{End}_A(M)$ -module Q_T is given by all homomorphisms from M to T , for abbreviation denoted by $Q_T = \text{Hom}_A(M, T) =: (M, T)$. To prove that $\text{gl.dim}(\text{End}_A(M)) \leq 3$, we explicitly construct a projective resolution with length ≤ 3 for every simple module E_T . The general method is as follows. Recall that for a finitely generated A -module M , $\text{add } M$ denotes the full subcategory of $\text{mod } A$ consisting of direct sums of direct summands of M . For any indecomposable summand T of M , we first construct a suitable exact sequence $0 \rightarrow K \rightarrow N_1 \rightarrow T$ with $N_1 \in \text{add } M$ with the following property:

(*) Every homomorphism from an indecomposable summand of M to T , except the multiples of the identity on T , factors through N_1 .

Applying the functor $(M, -)$, we get another short exact sequence

$$0 \rightarrow (M, K) \rightarrow (M, N_1) \rightarrow (M, T).$$

If the cokernel of $(M, N_1) \rightarrow (M, T)$ is 1-dimensional then the cokernel is E_T (because (M, T) is projective), i.e. we get the initial part of a projective resolution of the simple $\text{End}_A(M)$ -module E_T .

If $K \in \text{add } M$, then also (M, K) is projective and we have constructed a projective resolution of E_T of length 2. Otherwise, we construct another suitable short exact sequence $0 \rightarrow K' \rightarrow N_2 \rightarrow K$ with $N_2 \in \text{add } M$ with the following property:

(**) Every map from an indecomposable summand of M to K factors through N_2 .

Applying the functor $(M, -)$ to this short exact sequence, we get a short exact sequence

$$0 \rightarrow (M, K') \rightarrow (M, N_2) \rightarrow (M, K) \rightarrow 0.$$

If it happens that $K' \in \text{add } M$, we have (M, N_1) , (M, N_2) , (M, K') and (M, T) are all projective $\text{End}_A(M)$ -modules. Hence we get a projective resolution of the simple module E_T :

$$0 \rightarrow Q_{K'} \rightarrow Q_{N_2} \rightarrow Q_{N_1} \rightarrow Q_T \rightarrow E_T \rightarrow 0$$

and $\text{proj.dim } E_T \leq 3$.

The crucial aspect in this strategy is to come up with a generator-cogenerator M for which this process really stops at this stage, i.e. for which $K' \in \text{add } M$ in the second step above.

3 Proof of the main theorem

Before proving our main theorem, we introduce some notations. For any non-negative integer n , we use Λ_n to denote the (commutative) algebra $k[x, y]/(x^2, y^{n+2})$. The quotient algebra $\Lambda_n/\text{soc}(\Lambda_n)$ will be denoted by A_n .

Actually, in order to prove Theorem 1.1 it suffices to show that $\text{repdim}(A_n) = 3$, due to the following general result. Recall that an algebra A is called *basic* if all simple A -modules are of dimension 1. Note that our algebras $k[x, y]/(x^2, y^{2+n})$ are basic since the trivial module k is the only simple module.



Lemma 3.1 ([3], Proposition 1.2) *Let Λ be a basic algebra, and let P be an indecomposable projective-injective Λ -module. Define $A = \Lambda/\text{soc}(P)$. If $\text{rep.dim}(A) \leq 3$, then $\text{rep.dim}(\Lambda) \leq 3$.*

The structure of the projective-indecomposable A_n -module can be conveniently described diagrammatically as follows.



Here, the vertices correspond to basis vectors of the module and the edges describe the action of the generators x and y of A_n .

Now, we use a table to introduce and illustrate the notations for several classes of A_n -modules which will be used throughout the paper.

Notation	Shape of the A_n -module
$U_i, 0 \leq i \leq n+1$	
X	
$A_i^j, i, j \geq 0, i+j \leq n$	
$DA_i^j, i, j \geq 0, i+j \leq n$	

Note that the projective A_n -module will be denoted by A_n^0 . By Lemma 3.1, $\text{rep.dim}(A_n) \leq 3$ will imply that $\text{rep.dim}(\Lambda_n) \leq 3$. So we consider A_n instead of Λ_n . For any $n \geq 0$, we will prove that $\text{rep.dim}(A_n) \leq 3$ by constructing a generator-cogenerator M_n for A_n such that $\text{gl.dim}(\text{End}_{A_n}(M_n)) \leq 3$.

Now we give our constuction of the desired generator-cogenerators M_n of A_n as follows.

For $n = 0$ we put

$$M_0 = A_0^0 \oplus DA_0^0 \oplus U_0 \oplus U_1 \oplus X.$$

(Actually, for the algebra $A_0 = k[x, y]/(x^2, y^2, xy)$ it is known that $\text{repdim}(A_0) = 3$; in fact, A_0 is special biserial and we can apply Corollary 1.3 from [3]. However, we give the explicit generator-cogenerator to indicate the “induction base”.)

If n is positive, we put

$$M_n = \bigoplus_{i,j \geq 0, i+j \leq n} DA_i^j \oplus \bigoplus_{i \geq 0, j > 0, i+j \leq n} A_i^j \oplus A_n^0 \oplus \bigoplus_{0 \leq i \leq n+1} U_i \oplus X$$

Remark 3.2 Let us point out more clearly why this is an inductive construction, i.e. that one obtains M_n from M_{n-1} by a simple “combinatorial” procedure. Note that in M_n we still have all summands of M_{n-1} , except the projective module A_{n-1}^0 . The new summands in M_n are DA_i^{n-i} and A_i^{n-i} (for $0 \leq i \leq n$) and the uniserial module U_{n+1} . With the exception of A_n^0 and its dual, they can be obtained by extending from summands of M_{n-1} as follows. For all summands of M_{n-1} we add an additional vertex on top (i.e. an additional basis vector of the module) whenever possible; but we only allow this when no new squares are created in the shape of the module. All modules obtained in this way are added as summands of M_n (but only once, to avoid multiplicities).

For instance, going from M_0 to M_1 , the extension of DA_0^0 to the four-dimensional module having the shape of a square is not allowed; on the other hand, the extension of DA_0^0 to the four-dimensional module DA_0^1 is allowed and gives a new summand. With this procedure, for $i = 0, \dots, n-1$ the new modules DA_i^{n-i} are obtained from the summands DA_i^{n-i-1} of M_{n-1} , the new summands A_i^{n-i} are obtained from A_i^{n-i-1} and U_{n+1} is obtained from U_n . Note that the new projective module A_n^0 and the new injective module DA_n^0 can not be obtained with this method from summands of M_{n-1} , so they have to be added separately.

Hence, loosely speaking, the simple recipe for constructing M_n from M_{n-1} is as follows.

- (i) For all summands of M_{n-1} construct all possible modules obtained by adding a vertex on top, but without creating new squares in the shapes of the modules. Add these modules as new summands (if they are not already summands).
- (ii) Remove the old projective module A_{n-1}^0 . (But keep all other summands of M_{n-1} .)
- (iii) Add the new projective A_n^0 and its dual DA_n^0 as summands.

This inductive construction of a generator-cogenerator for A_n is actually the crucial step in the proof of our main result. We will show below that the endomorphism ring of M_n has global dimension 3, which then implies that A_n has representation dimension 3, as claimed.

We are now in the position to give the proof of our main result.

Proof of Theorem 1.1: We have to show that $\text{gl.dim}(\text{End}_{A_n}(M_n)) \leq 3$ for any $n \geq 0$.

(I) The case $n = 0$. We have $M_0 = A_0^0 \oplus DA_0^0 \oplus U_0 \oplus U_1 \oplus X$.

(1) The projective module A_0^0 . Since the radical of the projective module A_0^0 is $U_0 \oplus U_1 \in \text{add } M_0$ and the exact sequence $0 \rightarrow \text{rad } A_0^0 \rightarrow A_0^0$ clearly has the property (*), we can easily get a projective resolution of the simple $\text{End}_{A_0}(M_0)$ -module $E_{A_0^0}$:

$$0 \rightarrow Q_{U_0} \oplus Q_{U_1} \rightarrow Q_{A_0^0} \rightarrow E_{A_0^0} \rightarrow 0.$$

by applying the functor $(M_0, -)$ to $0 \rightarrow \text{rad } A_0^0 \rightarrow A_0^0$. Hence $\text{proj.dim } E_{A_0^0} = 1$.

(2) The injective module DA_0^0 . There is an exact sequence $0 \rightarrow U_0 \rightarrow U_1 \oplus X \rightarrow DA_0^0 \rightarrow 0$ with the property (*) (Note that there is no epimorphism from an indecomposable summand of M_0 to DA_0^0 , except the multiples of the identity on DA_0^0). Since U_0 and the middle term are in $\text{add } M_0$, by applying the functor $(M_0, -)$, we get a projective resolution of the simple $\text{End}_{A_0}(M_0)$ -module $E_{DA_0^0}$:

$$0 \rightarrow Q_{U_0} \rightarrow Q_{U_1} \oplus Q_X \rightarrow Q_{DA_0^0} \rightarrow E_{DA_0^0} \rightarrow 0.$$

Hence $\text{proj.dim } E_{DA_0^0} \leq 2$.

(3) The module X . There is a short exact sequence $0 \rightarrow U_0 \rightarrow A_0^0 \rightarrow X \rightarrow 0$ with the property (*) (Clearly, epimorphisms only occur from A_0^0 to X and they factor through the middle term. Those non-epimorphisms from an indecomposable summand of M_0 to X must have image in $\text{rad } X$ and clearly they also factor through the middle term.) Applying $(M_0, -)$ to the exact sequence, we get a projective resolution of the simple $\text{End}_{A_0}(M_0)$ -module E_X :

$$0 \rightarrow Q_{U_0} \rightarrow Q_{A_0^0} \rightarrow Q_X \rightarrow E_X \rightarrow 0$$

and hence $\text{proj.dim } E_X \leq 2$.

(4) The modules U_i where $i = 0, 1$. By symmetry, it suffices to show that $\text{proj.dim } E_{U_0} \leq 2$. For U_0 , there is a short exact sequence $0 \rightarrow K \rightarrow DA_0^0 \oplus DA_0^0 \rightarrow U_0 \rightarrow 0$ with the property (*) and the kernel K has the following shape

$$\begin{array}{c} \blacksquare \quad \blacktriangledown \quad \blacktriangledown \quad \blacksquare \\ \blacktriangledown \quad \blacksquare \quad \blacksquare \quad \blacktriangledown \end{array}$$

Since K is not a summand of M_0 , we need to find a short exact sequence ending at K with the property (**). There is a short exact sequence $0 \rightarrow U_0 \oplus U_0 \rightarrow U_1 \oplus A_0^0 \oplus X \rightarrow K \rightarrow 0$ with the property (**). Applying the functor $(M_0, -)$ to the above two short exact sequences and putting the resulting exact sequences together, we get a project resolution of E_{U_0} :

$$0 \rightarrow Q_{U_0} \oplus Q_{U_0} \rightarrow Q_{U_1} \oplus Q_{A_0^0} \oplus Q_X \rightarrow Q_{DA_0^0} \oplus Q_{DA_0^0} \rightarrow Q_{U_0} \rightarrow E_{U_0} \rightarrow 0$$

Hence $\text{proj.dim } E_{U_0} \leq 3$.

From the above, we have that $\text{gl.dim } (\text{End } M_0) \leq 3$, and as a consequence $\text{reproj.dim } (A_0) \leq 3$. Since A_0 is not of finite representation type we can actually deduce that $\text{reproj.dim } (A_0) = 3$.

(II) The case $n > 0$. We have

$$M_n = \bigoplus_{i,j \geq 0, i+j \leq n} DA_i^j \oplus \bigoplus_{i \geq 0, j > 0, i+j \leq n} A_i^j \oplus A_n^0 \oplus \bigoplus_{0 \leq i \leq n+1} U_i \oplus X.$$

Again, for any indecomposable summand N of M_n , we show that $\text{proj.dim } E_N \leq 3$ by explicitly constructing a projective resolution for the corresponding simple $\text{End}_{A_n}(M_n)$ -module E_N . For the convenience of the reader, we list the indecomposable summands of M_n as follows.

$$\begin{array}{cccccc} A_n^0 & & & & & X \\ & A_{n-1}^1 & A_{n-2}^1 & \cdots & A_1^1 & A_0^1 \\ & & A_{n-2}^2 & \cdots & A_1^2 & A_0^2 \\ & & & \ddots & \vdots & \vdots \\ & & & & A_1^{n-1} & A_0^{n-1} \\ & & & & & A_0^n \end{array}$$

$$\begin{array}{cccccc}
DA_n^0 & DA_{n-1}^0 & DA_{n-2}^0 & \cdots & DA_1^0 & DA_0^0 \\
& DA_{n-1}^1 & DA_{n-2}^1 & \cdots & DA_1^1 & DA_0^1 \\
& & DA_{n-2}^2 & \cdots & DA_1^2 & DA_0^2 \\
& & & \ddots & \vdots & \vdots \\
& & & & DA_1^{n-1} & DA_0^{n-1} \\
& & & & & DA_0^n
\end{array}$$

$$U_0, U_1, U_2, \dots, U_n, U_{n+1}$$

(1) The projective module A_n^0 . Clearly, the exact sequence $0 \rightarrow \text{rad } A_n^0 \rightarrow A_n^0 \rightarrow 0$ has the property (*). Since $\text{rad } A_n^0 = A_{n-1}^1 \in \text{add } M_n$, applying the functor $(M_n, -)$ to the exact sequence gives us a projective resolution of the simple $\text{End}_{A_n}(M_n)$ module $E_{A_n^0}$ of the form

$$0 \rightarrow Q_{A_{n-1}^1} \rightarrow Q_{A_n^0} \rightarrow E_{A_n^0} \rightarrow 0.$$

Hence $\text{proj.dim } E_{A_n^0} = 1$.

(2) The module X . There is a short exact sequence $0 \rightarrow A_{n-1}^1 \rightarrow U_0 \oplus A_n^0 \rightarrow X \rightarrow 0$ with the property (*) (Actually, the only epimorphisms, apart from the multiples of the identity on X , are from A_n^0 , so clearly they factor through the middle term. All other maps factor through the radical $\text{rad } X = U_0$). Since A_{n-1}^1 and the middle term are in $\text{add } M_n$, applying the functor $(M_n, -)$, we get a projective resolution of the simple $\text{End}_{A_n}(M_n)$ -module E_X :

$$0 \rightarrow Q_{A_{n-1}^1} \rightarrow Q_{U_0} \oplus Q_{A_n^0} \rightarrow Q_X \rightarrow E_X \rightarrow 0$$

Hence $\text{proj.dim } E_X \leq 2$.

(3) The modules A_i^1 . If $i = 0$, then there is a short exact sequence $0 \rightarrow DA_0^1 \rightarrow U_1 \oplus U_0 \oplus DA_1^0 \rightarrow A_0^1 \rightarrow 0$ with the property (*) (Except for the multiples of the identity on A_0^1 , the epimorphisms are from A_s^1 ($s > 0$) and DA_t^0 . They all factor through DA_1^0). Since all terms of the exact sequence are in $\text{add } M_n$, by applying the functor $(M_n, -)$, we get a projective resolution of the simple $\text{End}_{A_n}(M_n)$ -module $E_{A_0^1}$:


$$0 \rightarrow Q_{DA_0^1} \rightarrow Q_{U_1} \oplus Q_{U_0} \oplus Q_{DA_1^0} \rightarrow Q_{A_0^1} \rightarrow E_{A_0^1} \rightarrow 0$$

Hence $\text{proj.dim } E_{A_0^1} \leq 2$.

If $i > 0$, then there is a short exact sequence $0 \rightarrow DA_i^1 \rightarrow A_{i-2}^2 \oplus DA_{i+1}^0 \rightarrow A_i^1$ with the property (*) (Apart from the multiples of the identity on A_i^1 , the epimorphisms to A_i^1 are from A_s^1 ($s > i$) and DA_t^1 ($t > i$). They all factor through DA_i^1 . The non-epimorphisms all factor through the middle term by the definition of the middle term). Note that all the terms of the short exact sequence are in $\text{add } M_n$. Applying the functor $(M_n, -)$, we get a projective resolution of the simple $\text{End}_{A_n}(M_n)$ -module $E_{A_i^1}$:

$$0 \rightarrow Q_{DA_i^1} \rightarrow Q_{A_{i-1}^2} \oplus Q_{DA_{i+1}^0} \rightarrow Q_{A_i^1} \rightarrow E_{A_i^1} \rightarrow 0$$

(4) The modules A_i^j ($j > 1, i > 0$). For every such module, there is a short exact sequence $0 \rightarrow K \rightarrow A_{i-1}^{j+1} \oplus A_i^{j-1} \oplus DA_{i+1}^{j-1} \rightarrow A_i^j \rightarrow 0$ with the property (*) (In fact, the only epimorphisms from summands other than A_i^j are from A_s^j ($s > i$) and DA_t^{j-1} ($t > i$). They all factor through DA_{i+1}^{j-1} . Other maps factor through $A_{i-1}^{j+1} \oplus A_i^{j-1}$). The kernel K has the following shape:




Clearly, K is not in $\text{add } M_n$. There is a short exact sequence $0 \rightarrow DA_i^{j-1} \rightarrow A_{i-1}^j \oplus DA_i^j \oplus DA_{i+1}^{j-2} \rightarrow K \rightarrow 0$ with the property (**). Applying the functor $(M_n, -)$ to the above two exact sequences and putting the resulting sequences together, we get a projective resolution of the simple $\text{End}_{A_n}(M_n)$ -module $E_{A_i^j}$:

$$0 \rightarrow Q_{DA_i^{j-1}} \rightarrow Q_{A_{i-1}^j} \oplus Q_{DA_i^j} \oplus Q_{DA_{i+1}^{j-2}} \rightarrow Q_{A_{i-1}^{j+1}} \oplus Q_{A_i^{j-1}} \oplus Q_{DA_{i+1}^{j-1}} \rightarrow Q_{A_i^j} \rightarrow E_{A_i^j} \rightarrow 0$$

Hence $\text{proj.dim } E_{A_i^j} \leq 3$ for $j > 1$ and $i > 0$.

(5) The modules A_0^j ($j > 1$). For any A_0^j with $j > 1$, there is a short exact sequence $0 \rightarrow K \rightarrow U_j \oplus A_0^{j-1} \oplus DA_1^{j-1} \rightarrow A_0^j \rightarrow 0$ with the property (*) (The only epimorphisms not from A_0^j itself are from A_s^j ($s > 0$) or DA_t^j ($t > 0$). They all factor through DA_1^{j-1}). The kernel K has the following shape:



Again, K is not in $\text{add } M_n$, but there is a short exact sequence $0 \rightarrow U_j \rightarrow U_{j-1} \oplus U_{j+1} \oplus DA_1^{j-2} \rightarrow K \rightarrow 0$ with the property (**). Applying the functor $(M_n, -)$ to the above two short exact sequences and putting together the resulting sequences, we get a projective resolution of the simple $\text{End}_{A_n}(M_n)$ -module $E_{A_0^j}$:

$$0 \rightarrow Q_{U_j} \rightarrow Q_{U_{j-1}} \oplus Q_{U_{j+1}} \oplus Q_{DA_1^{j-2}} \rightarrow Q_{U_j} \oplus Q_{A_0^{j-1}} \oplus Q_{DA_1^{j-1}} \rightarrow Q_{A_0^j} \rightarrow E_{A_0^j} \rightarrow 0$$

Hence $\text{proj.dim } E_{A_0^j} \leq 3$ for $j > 1$.

(6) The module DA_n^0 . There is a short exact sequence $0 \rightarrow A_{n-1}^1 \rightarrow DA_{n-1}^1 \oplus A_n^0 \rightarrow DA_n^0 \rightarrow 0$ with the property (*) and all terms of the short exact sequence are in $\text{add } M_n$. Applying the functor $(M_n, -)$ to the short exact sequence, we get a projective resolution of the simple $\text{End}_{A_n}(M_n)$ -module $E_{DA_n^0}$:

$$0 \rightarrow Q_{A_{n-1}^1} \rightarrow Q_{DA_{n-1}^1} \oplus Q_{A_n^0} \rightarrow Q_{DA_n^0} \rightarrow E_{DA_n^0} \rightarrow 0$$

Hence $\text{proj.dim } DA_n^0 \leq 2$.

(7) The modules $DA_i^0 (0 < i < n)$. For any DA_i^0 with $0 < i < n$, there is a short exact sequence $0 \rightarrow A_{i-1}^2 \rightarrow DA_{i-1}^1 \oplus A_i^1 \rightarrow DA_i^0 \rightarrow 0$ with the property (*) (Except for the multiples of the identity on DA_i^0 , the only epimorphisms to DA_i^0 are from $DA_s^0 (s > i)$ and $A_t^1 (t \geq i)$. They all factor through A_i^1). Note that all terms of the short exact sequence are in $\text{add } M_n$. Applying the functor $(M_n, -)$, we get a projective resolution of $E_{DA_i^0}$:

$$0 \rightarrow Q_{A_{i-1}^2} \rightarrow Q_{DA_{i-1}^1} \oplus Q_{A_i^1} \rightarrow Q_{DA_i^0} \rightarrow E_{DA_i^0} \rightarrow 0$$

Hence we have $\text{proj.dim } E_{DA_i^0} \leq 2$ for $0 < i < n$.

(8) The module DA_0^0 . There is a short exact sequence $0 \rightarrow K \rightarrow X \oplus A_0^1 \rightarrow DA_0^0 \rightarrow 0$ with the property (*) (Note that all epimorphisms from an indecomposable summand of M_n to DA_0^0 , except the multiples of the identity on DA_0^0 , factor through A_0^1). The kernel K is isomorphic to A_0^0 , which is not in $\text{add } M_n$. There is a short exact sequence $0 \rightarrow A_{n-1}^1 \rightarrow U_0 \oplus U_0 \oplus A_n^0 \rightarrow K \rightarrow 0$ with the property (**). Applying the functor $(M_n, -)$ to the above two exact sequences and putting the resulting sequences together, we get a projective resolution of the simple $\text{End } M_n$ module $E - DA_0^0$:

$$0 \rightarrow Q_{A_{n-1}^1} \rightarrow Q_{U_0} \oplus Q_{U_0} \oplus Q_{A_n^0} \rightarrow Q_X \oplus Q_{A_0^1} \rightarrow Q_{DA_0^0} \rightarrow E_{DA_0^0} \rightarrow 0$$

Hence we have $\text{proj.dim } E_{DA_0^0} \leq 3$.

(9) The modules $DA_i^j (i, j > 0, i+j = n)$. For each module DA_i^j with $i, j > 0$ and $i+j = n$, there is a short exact sequence $0 \rightarrow DA_{i-1}^j \rightarrow DA_{i-1}^{j+1} \oplus DA_i^{j-1} \rightarrow DA_i^j$ with the property (*) and all the terms of the short exact sequence are in $\text{add } M_n$. Applying the functor $(M_n, -)$ to the exact sequence, we get a projective resolution of the simple $\text{End } M_n$ module $E_{DA_i^j}$:

$$0 \rightarrow Q_{DA_{i-1}^j} \rightarrow Q_{DA_{i-1}^{j+1}} \oplus Q_{DA_i^{j-1}} \rightarrow Q_{DA_i^j} \rightarrow E_{DA_i^j} \rightarrow 0$$

Hence we have $\text{proj.dim } E_{DA_i^j} \leq 2$ for $i, j > 0$ and $i+j = n$.

(10) The modules $DA_0^j (j > 0)$. First, we consider DA_0^n . There is a short exact sequence $0 \rightarrow U_n \rightarrow U_{n+1} \oplus DA_0^{n-1} \rightarrow DA_0^n$ with the property (*). Applying the functor $(M_n, -)$ to the short exact sequence, we get a projective resolution of the simple $\text{End } A_n(M_n)$ -module $E_{DA_0^n}$:


$$0 \rightarrow Q_{U_n} \rightarrow Q_{U_{n+1}} \oplus Q_{DA_0^{n-1}} \rightarrow Q_{DA_0^n} \rightarrow E_{DA_0^n} \rightarrow 0$$

Hence we have $\text{proj.dim } E_{DA_0^n} \leq 2$. For each DA_0^j with $0 < j < n$, there is a short exact sequence $0 \rightarrow A_0^j \rightarrow DA_0^{j-1} \oplus A_0^{j+1} \rightarrow DA_0^j \rightarrow 0$ with the property (*). Applying the functor $(M_n, -)$ to the short exact sequence, we get a projective resolution of the simple $\text{End } A_n(M_n)$ -module $E_{DA_0^j}$:

$$0 \rightarrow Q_{A_0^j} \rightarrow Q_{DA_0^{j-1}} \oplus Q_{A_0^{j+1}} \rightarrow Q_{DA_0^j} \rightarrow E_{DA_0^j} \rightarrow 0$$

Hence we have $\text{proj.dim } E_{DA_0^j} \leq 2$.

(11) The modules $DA_i^j(i, j > 0, i + j < n)$. For each module DA_i^j with $i, j > 0$ and $i + j < n$, there is a short exact sequence $0 \rightarrow K \rightarrow DA_i^{j-1} \oplus DA_{i-1}^{j+1} \oplus A_i^{j+1} \rightarrow DA_i^j$ with the property (*) (Except the multiples of the identity on DA_i^j , the only epimorphisms to DA_i^j are from DA_s^j ($s > i$) and A_t^{j+1} ($t \geq i$). They all factor through A_i^{j+1} . Other maps factor through the middle term by the definition of the middle term.) The kernel K has the following shape:




Clearly, K is not in $\text{add } M_n$. However, there is a short exact sequence $0 \rightarrow A_{i-1}^{j+1} \rightarrow DA_{i-1}^j \oplus A_i^j \oplus A_{i-1}^{j+2} \rightarrow K \rightarrow 0$ with the property (**). Applying the functor $(M_n, -)$ to the above two exact sequences and putting the resulting sequences together, we get a projective resolution of the simple $\text{End}_{A_n}(M_n)$ -module $E_{DA_i^j}$:

$$0 \rightarrow Q_{A_{i-1}^{j+1}} \rightarrow Q_{DA_{i-1}^j} \oplus Q_{A_i^j} \oplus Q_{A_{i-1}^{j+2}} \rightarrow Q_{DA_{i-1}^{j-1}} \oplus Q_{DA_{i-1}^{j+1}} \oplus Q_{A_i^{j+1}} \rightarrow Q_{DA_i^j} \rightarrow E_{DA_i^j} \rightarrow 0$$

Hence we have $\text{proj.dim } E_{DA_i^j} \leq 3$ for $i, j > 0$ and $i + j < n$.

(12) The module U_0 . There is a short exact sequence $0 \rightarrow K \rightarrow DA_0^n \oplus DA_0 \rightarrow U_0 \rightarrow 0$ with the property (*) (Except the multiples of the identity on U_0 , all maps from an indecomposable summand of M_n to U_0 factor either through DA_0^n or DA_0). The kernel K has the following shape:




Clearly, K is not in $\text{add } M_n$. However, there is a short exact sequence $0 \rightarrow U_n \oplus U_0 \rightarrow U_{n+1} \oplus A_n^0 \oplus X \rightarrow K \rightarrow 0$ with the property (**). Applying the functor $(M_n, -)$ to the above exact sequence and putting the resulting sequences together, we get a projective resolution of the simple $\text{End}_{A_n}(M_n)$ -module E_{U_0} :

$$0 \rightarrow Q_{U_n} \oplus Q_{U_0} \rightarrow Q_{U_{n+1}} \oplus Q_{A_0^n} \oplus Q_X \rightarrow Q_{DA_0^n} \oplus Q_{DA_0} \rightarrow Q_{U_0} \rightarrow E_{U_0} \rightarrow 0$$

Hence we have $\text{proj.dim } E_{U_0} \leq 3$.

(13) The module U_1 . There is a short exact sequence $0 \rightarrow K \rightarrow DA_0^1 \oplus A_0^n \rightarrow U_1 \rightarrow 0$ with the property (*) (Except the multiples of the identity on U_1 , every epimorphism from an indecomposable summand of M_n to U_1 factors through either DA_0^1 or A_0^n . The maps having image in the radical of U_1 factor through A_0^n). The kernel K has the following shape:




There is another short exact sequence $0 \rightarrow A_0^n \rightarrow U_n \oplus A_1^{n-1} \oplus DA_0^0 \rightarrow K \rightarrow 0$ with the property (**). Applying the functor $(M_n, -)$ to the above two exact sequences and putting the resulting sequences together, we get a projective resolution of the simple $\text{End}_{A_n}(M_n)$ -module E_{U_1} :

$$0 \rightarrow Q_{A_0^n} \rightarrow Q_{U_n} \oplus Q_{A_1^{n-1}} \oplus Q_{DA_0^0} \rightarrow Q_{DA_0^1} \oplus Q_{A_0^n} \rightarrow Q_{U_1} \rightarrow E_{U_1} \rightarrow 0$$

Hence we have $\text{proj.dim } E_{U_1} \leq 3$.

(14) The modules U_i , $(1 < i < n+1)$. For each module U_i with $1 < i < n+1$, there is a short exact sequence $0 \rightarrow K \rightarrow U_{i-1} \oplus DA_0^i \oplus A_{i-1}^{n-i+1} \rightarrow U_i \rightarrow 0$ with the property (*) (In fact, the epimorphisms from an indecomposable summand of M_n to U_i , except the multiples of the identity on U_i , either factor through DA_0^i or A_{i-1}^{n-i+1} . Other maps factor through the U_{i-1} , which is the radical of U_i). The kernel K has the following shape:



There is a short exact sequence $0 \rightarrow A_{i-1}^{n-i+1} \xrightarrow{2} A_i^{n-i} \oplus A_{i-2}^{n-i+2} \oplus DA_0^{i-1} \rightarrow K \rightarrow 0$ with the property (**). Applying the functor $(M_n, -)$ to the above two exact sequences and putting together the resulting sequences, we get a projective resolution of the simple $\text{End } M_n$ module E_{U_i} :

$$0 \rightarrow Q_{A_{i-1}^{n-i+1}} \rightarrow Q_{A_i^{n-i}} \oplus Q_{A_{i-2}^{n-i+2}} \oplus Q_{DA_0^{i-1}} \rightarrow Q_{U_{i-1}} \oplus Q_{DA_0^i} \oplus Q_{A_{i-1}^{n-i+1}} \rightarrow Q_{U_i} \rightarrow E_{U_i} \rightarrow 0$$

Hence $\text{proj.dim } E_{U_i} \leq 3$ for $1 < i < n+1$.

(15) The module U_{n+1} . There is a short exact sequence $0 \rightarrow A_{n-1}^1 \rightarrow U_n \oplus A_n^0 \rightarrow U_{n+1} \rightarrow 0$ with the property (**). (In fact, except the multiples of the identity on U_{n+1} , the only epimorphisms are from A_n^0 . They factor through A_n^0 . Other maps factor through the radical U_n). Since all terms of the short exact sequence are in $\text{add } M_n$, by applying the functor $(M_n, -)$, we get a projective resolution of the simple $\text{End}_{A_n}(M_n)$ -module $E_{U_{n+1}}$:

$$0 \rightarrow Q_{A_{n-1}^1} \rightarrow Q_{U_n} \oplus Q_{A_n^0} \rightarrow Q_{U_{n+1}} \rightarrow E_{U_{n+1}} \rightarrow 0$$

Hence we have $\text{proj.dim } E_{U_{n+1}} \leq 2$.

Thus we have shown that $\text{gl.dim } \text{End}_{A_n}(M_n) \leq 3$ for any non-negative integer n . Hence we have $\text{rep.dim } A_n \leq 3$ for all $n \geq 0$. On the other hand, the algebras A_n ($n \geq 0$) are not representation finite, so we have $\text{rep.dim } A_n > 2$ for all $n \geq 0$. Altogether, we come to the conclusion that $\text{rep.dim } A_n = 3$ for all non-negative integers n and this completes the proof of Theorem 1.1. $\diamond$

References

- [1] M. AUSLANDER, Representation dimension of Artin algebras. Queen Mary College, Mathematics Notes, University of London. 1971. Also in: I. Reiten, S. Smalø, Ø. Solberg (Eds.), Selected works of Maurice Auslander, Part I, Amer. Math. Soc., Providence, RI, 1999, pp. 505-574.
- [2] V.M. BONDARENKO, J.A. DROZD, The representation type of finite groups. (Russian) Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) 71 (1977), 24-41. English translation: J. Soviet Math. 20 (1982), 2515-2528.
- [3] K. ERDMANN, T. HOLM, O. IYAMA AND J. SCHRÖER, Radical embeddings and representation dimension. Advances in Math. 185 (2004), 159-177.
- [4] X. GUO, Representation dimension: An invariant under stable equivalence. Trans. Amer. Math. Soc. 357 (2005) 3255-3263.
- [5] T. HOLM, Representation dimension of some tame blocks of finite groups. Algebra Colloq. 10 (2003), 275-284.
- [6] O. IYAMA, Finiteness of representation dimension. Proc. Amer. Math. Soc. 131 (2003), 1011-1014.
- [7] O. IYAMA, Rejective subcategories of artin algebras and orders. Preprint (2003), math.RT/0311281.
- [8] R. ROUQUIER, Dimensions of triangulated categories. Preprint (2003), math.CT/0310134
- [9] R. ROUQUIER, Representation dimension of exterior algebras. Preprint (2005), available at <http://www.math.jussieu.fr/~rouquier/preprints/preprints.html>
- [10] C. XI, Representation dimension and quasi-hereditary algebras. Advances in Math. 168 (2002), 193-212.

THORSTEN HOLM

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF LEEDS
LEEDS LS2 9JT, ENGLAND
E-mail: tholm@maths.leeds.ac.uk

WEI HU

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF LEEDS
LEEDS LS2 9JT, ENGLAND
AND

SCHOOL OF MATHEMATICAL SCIENCES, BEIJING NORMAL UNIVERSITY
100875 BEIJING, P. R. CHINA
E-mail: huwei@maths.leeds.ac.uk